

Appendix 2. Prey types used by Mountain Bluebird (*Sialia currucoides*) parents to feed their nestlings.

Table A2.1. The percentage of all prey types that were fed to nestlings in clear-cuts and grasslands. Cicadas were separated from Hemiptera because their bodies were relatively large and bulky. Unknown refers to (usually small), unidentifiable prey items or cases where parents brought multiple items that could not be distinguished. There were 2205 prey items brought in clear-cuts and 2662 prey items brought in grasslands. Prey types are roughly ordered in decreasing abundance.

Prey Type	Clear-cuts	Grasslands
Larva	40.54%	28.21%
Spider	26.44%	15.18%
Coleoptera	9.84%	25.21%
Unknown	6.53%	11.46%
Orthoptera	5.49%	6.31%
Hymenoptera	4.63%	4.40%
Lepidoptera	3.40%	2.82%
Diptera	1.77%	2.52%
Odonata	0.63%	1.54%
Cicada	0.18%	1.43%
Ephemeroptera	0.09%	0.41%
Trichoptera	0.09%	0.3%
Annelida	0.27%	0.04%
Hemiptera	0.09%	0.04%
Neuroptera	0%	0.08%
Myriapoda	0%	0.04%
Plecoptera	0%	0.04%