

Appendix 2. Relationships between landscape variables (in terms of area or percentage within the home range) and the average number of Pileated Woodpecker chicks fledged between 2009 and 2013 in the study area.

Variable	Parameter			Regression		<i>p</i>
	Estimate	Std Error	<i>t</i>	Adj R-sq	<i>F</i> <sub>1,7</sub>	
<b>Area</b>						
Coniferous	0.0006	0.006	0.1	-0.14	0.01	0.92
Deciduos/mixed	0.0035	0.008	0.42	-0.12	0.17	0.69
Regeneration forest	0.0061	0.029	0.21	-0.14	0.04	0.84
Grassland	0.0224	0.018	1.24	0.06	1.55	0.25
Light Urban	0.0056	0.005	1.03	0.01	1.06	0.34
Medium Urban	-0.0019	0.007	-0.3	-0.13	0.09	0.77
Heavy Urban	-0.0016	0.022	-0.07	-0.14	0.01	0.95
Other	-0.0889	0.05	-1.78	0.21	3.16	0.12
<b>Percentage</b>						
Coniferous	0.0051	0.021	0.24	-0.13	0.57	0.82
Deciduos/mixed	-0.0072	0.041	-0.18	-0.14	0.03	0.87
Regeneration forest	-0.0463	0.096	-0.48	-0.11	0.23	0.65
Grassland	0.1087	0.078	1.4	0.11	1.95	0.21
Light Urban	0.029	0.027	1.07	0.02	1.15	0.32
Medium Urban	-0.0106	0.017	-0.62	-0.08	0.39	0.55
Heavy Urban	-0.0215	0.078	-0.28	-0.13	0.08	0.79
Other	-0.3254	0.714	-0.46	-0.11	0.21	0.66