

Appendix 1. Total birds observed in the four sampling sites (control, “burned 2002”, “burned 2005” and “burned 2002-2015”) during the 2016-2017 and 2017-2018 breeding seasons.

Name	Control	Burned 2002	Burned 2015	Burned 2002-2015
<i>Patagioenas araucana</i>	X			
<i>Enicognathus ferrugineus</i>	X	X	X	X
<i>Sephanoides sephaniodes</i>	X	X	X	
<i>Veniliornis lignarius</i>	X	X		
<i>Colaptes pitius</i>	X	X	X	
<i>Campephilus magellanicus</i>	X	X		
<i>Aphrastura spinicauda</i>	X	X	X	
<i>Sylviorthorhynchus desmursii</i>	X	X		X
<i>Pygarrhichas albogularis</i>	X	X	X	
<i>Pteroptochos tarnii</i>	X	X		
<i>Scelorchilus rubecola</i>	X	X	X	
<i>Scytalopus magellanicus</i>	X	X	X	X
<i>Elaenia albiceps</i>	X	X	X	X
<i>Xolmis pyrope</i>	X	X	X	X
<i>Colorhamphus parvirostris</i>	X			
<i>Tachycineta meyeri</i>	X	X	X	X
<i>Troglodytes aedon</i>	X	X	X	X
<i>Turdus falckandii</i>	X	X	X	X
<i>Phrygilus patagonicus</i>	X	X	X	X
<i>Spinus barbata</i>	X	X	X	X
<i>Glaucidium nanum</i>			X	
<i>Cinclodes fuscus</i>			X	
<i>Cinclodes patagonicus</i>			X	X
<i>Callipepla californica</i>			X	
<i>Zonotrichia capensis</i>		X	X	X
<i>Geranoaetus polyosoma</i>		X		
<i>Anairetes parulus</i>		X		X
<i>Notiochelidon cyanoleuca</i>		X		X
<i>Upecerthia dumetaria</i>		X		X
<i>Megaceryle torquata</i>				X
<i>Phytotoma rara</i>				X
<i>Sicalis luteola</i>				X
<i>Phrygilus fruticeti</i>				X
<i>Agriornis livida</i>				X
<i>Leptasthenura aegithaloides</i>		X		X