

**Appendix 7.** Long-term assessment of a published hypothesis of 11 extirpated bird species in the Cauca valley, using additional records after Restrepo and Naranjo (1987).

Family	Species	Number of Records	Records	Long-Term Diagnostic after 1987
Anatidae	<i>Cairina moschata</i>	7	1995, 2004, 2006, 2008, 2011, 2018, 2019	Not extirpated
Anatidae	<i>Spatula clypeata</i>	3	1995, 2017, 2020	Sporadic
Anatidae	<i>Mareca americana</i>	1	2014	Rare, Erratic
Anatidae	<i>Anas acuta</i>	3	2012, 2013, 2017, 2018	Local, Erratic
Anatidae	<i>Anas georgica</i>	5	2012, 2013, 2014, 2015, 2017, 2019	Recolonizer
Anatidae	<i>Aythya affinis</i>	1	2016	Rare, Sporadic
Anatidae	<i>Netta erythrophthalma</i>	0	NA	Extirpated
Ciconiidae	<i>Jabiru mycteria</i>	0	NA	Extirpated
Ciconiidae	<i>Mycteria americana</i>	2	2005, 2012	Extirpated
Ardeidae	<i>Tigrisoma lineatum</i>	3	1992, 2018, 2019	Recolonizer
Threskiornithidae	<i>Eudocimus albus</i>	0	NA	Extirpated

Restrepo, C. and L. G. Naranjo. 1987. Recuento histórico de la disminución de humedales y la desaparición de aves acuáticas en el valle geográfico del Río Cauca, Colombia. In: Memorias III Congreso de Ornitología Neotropical, Cali, Colombia, pp. 43–45.