

Appendix 4. Summary statistics for stable hydrogen isotope ($\delta^2\text{H}$) of insects separated by taxa, and site. Insect samples analyzed were collected from June 11 to July 16, 2020, and were homogenized and pooled to obtain the target mass necessary for hydrogen isotope analysis.

Site	Coleoptera $\delta^2\text{H}$ (<i>mean</i> \pm <i>SD</i> ‰)	n	Nematocera $\delta^2\text{H}$ (<i>mean</i> \pm <i>SD</i> ‰)	n
SSP01	-183.8 \pm 3.5	4	-167.6 \pm 5.0	6
SSP02	-213.0 \pm 22.2	6	-181.1 \pm 6.5	5
SSP03	-159.6 \pm 5.3	6	-180.0 \pm 4.0	6
SSP04	-218.4 \pm 41.9	6	-176.1 \pm 9.6	2
SSP05	-188.7 \pm 4.6	8	-189.2 \pm 16.9	6
SSP07	-183.2 \pm 16.3	6	-176.3 \pm 11.1	6
SSP08	-181.6 \pm 2.0	6	-216.4 \pm 5.4	6
SSP09	-202.2 \pm 22.3	4	-184.2 \pm 18.5	7
SSP10	-181.2 \pm 13.0	6	-197.6 \pm 8.9	7
SSP11	-177.8 \pm 7.6	6	-180.9 \pm 7.3	7
SSP12	-169.7 \pm 6.5	6	-184.6 \pm 5.3	6
SSP13	-205.3 \pm 4.9	6	-180.6 \pm 5.1	6
General	-185.18 \pm 14.82	70	-186 \pm 19.58	70